

$S_{\text{eff}}(E)$ and the ${}^7\text{Be}(p, \gamma){}^8\text{B}$ reaction

B. K. Jennings and S. Karataglidis

TRIUMF, 4004 Wesbrook Mall, Vancouver, British Columbia, Canada, V6T 2A3

(February 9, 2008)

Abstract

We explore approximations to the effective S factor for the ${}^7\text{Be}(p, \gamma){}^8\text{B}$ reaction using different approximation for the integral over the Gamov peak. In the temperature range of interest for solar neutrino production, $S_{\text{eff}}(E)$ may be determined to within 0.5% from $S(20)$ with no direct information on the derivatives of $S(E)$.

The ${}^7\text{Be}(p, \gamma){}^8\text{B}$ reaction, at energies of approximately 20 keV, plays an important role in the production of solar neutrinos [1]. The subsequent decay of the ${}^8\text{B}$ is the source of the high energy neutrinos to which many solar neutrino detectors are sensitive. The cross section for this reaction is conventionally expressed in terms of the S factor which is defined in terms of the cross section, σ , by

$$S(E) = \sigma(E)E \exp[2\pi\eta(E)] , \quad (1)$$

where $\eta(E) = Z_1 Z_2 \alpha \sqrt{\mu c^2 / 2E}$ is the Sommerfeld parameter, α is the fine structure constant, and μ is the reduced mass. The reaction rate per particle pair is [2]

$$\langle \sigma \nu \rangle = \left(\frac{8}{\pi \mu} \right)^2 \frac{1}{(kT)^{3/2}} \int_0^\infty dE S(E) \exp \left[-\frac{E}{kT} - \frac{b}{E^{1/2}} \right] \quad (2)$$

where T is the temperature, k is the Boltzmann factor and $b = (2\mu)^{1/2} \pi e^2 Z_1 Z_2 / \hbar$ comes from the Coulomb penetration factor. In order to study the approximations to the integral in Eq. (2) it is convenient to introduce an effective S factor defined by [2]

$$S_{\text{eff}}(E) = \exp[\tau] E_0 \sqrt{\frac{\tau}{4\pi}} \int_0^\infty dE S(E) \exp \left[-\frac{E}{kT} - \frac{b}{E^{1/2}} \right] \quad (3)$$

where $\tau = 3E_0/kT$ and $E_0 = (bkT/2)^{2/3}$ is the location at which the function $\exp \left[-\frac{E}{kT} - \frac{b}{E^{1/2}} \right]$ peaks. Since the integrand is strongly peaked the integral may be done by the saddle point method [2]. To first order in $1/\tau$ the result is [2,3]

$$S_{\text{eff-S}}(E) = S(E_0) \left\{ 1 + \frac{1}{\tau} \left[\frac{5}{12} + \frac{5S'(E_0)E_0}{S(E_0)} + \frac{S''(E_0)E_0^2}{S(E_0)} \right] \right\} . \quad (4)$$

The approximation depends on S and its first two derivatives, all evaluated at $E = E_0$. By expanding S about $E = 0$ MeV we obtain an expression where we only need S and its derivatives at the origin. This results in the approximation [3]

$$S_{\text{eff-T}} = S(0) \left[1 + \frac{5}{12\tau} + \frac{S'(0)(E_0 + \frac{35}{36}kT)}{S(0)} + \frac{S''(0)E_0}{S(0)} \left(\frac{E_0}{2} + \frac{89}{72}kT \right) \right] . \quad (5)$$

Eqs. (4) and (5) both explicitly depend on the first and second derivatives of the S factor. Another approximation may be made in which the derivatives in Eq. (4) are neglected, giving a much simpler form:

$$S_{\text{eff-Snd}} = S(E_0) \left\{ 1 + \frac{5}{12\tau} \right\} . \quad (6)$$

Finally we may replace $S(E_0)$ with $S(20)$ in this last equation to yield

$$S_{\text{eff-S20}} = S(20) \left\{ 1 + \frac{5}{12\tau} \right\} . \quad (7)$$

To evaluate these approximations we need a functional form for $S(E)$. For $E < 100$ keV, the functional form of S for the ${}^7\text{Be}(p, \gamma)$ reaction is well approximated by [4]

$$S(E)/S(0) = \frac{0.0409}{0.1375 + E} + 0.703 + 0.343E \quad (8)$$

where E is in MeV. With this expression for $S(E)/S(0)$ and taking $S(0) = 19.0$ eVb [4], we have calculated S_{eff} and the different approximations to it. The results are given in Table I for a range of temperatures that includes those relevant to the production of ${}^8\text{B}$ in the sun, i.e. $1.3 \times 10^7 \text{ K} < T < 1.6 \times 10^7 \text{ K}$ [3]. The values of S_{eff} calculated with any of the approximations deviate from the exact results, Eq. (3), by no more than 0.1 eVb or 0.5%. Also, there is less than a 0.1 eVb variation in S_{eff} over the given range of temperatures. We note particularly the accuracy of Eq. (7), which requires only the value of S at 20 keV, and does not require any explicit knowledge of the derivatives or of the temperature dependence in E_0 .

Since the first and second derivatives of the S factor enter into two of the approximations, an understanding of the accuracy with which these derivatives can be determined is useful. In Table II, we show the derivatives given in Ref. [4] for a range of models. The first four are hard sphere models [4] depending on the given hard sphere radius, r_c . The next three are Woods-Saxon models. These models agree with generator coordinate model calculations for low energies, $E < 300$ keV [4], so the conclusions drawn here will also apply to results from the generator coordinate model calculations.

The first derivative shows more model dependence than the second derivative. However even for the first derivative the model dependence is quite small. Neglecting the hard sphere model with $r_c = 4.1$ fm, which was shown to be unrealistic [4], and the results from Adelberger *et al.* [3], the average of the values of the first derivative is $(1/S)(dS/dE)|_{E=0} = -1.87 \pm 0.1 \text{ MeV}^{-1}$. This introduces a variation of 0.2% in S_{eff} and is negligible.

There is surprisingly little variation in the second derivative. In the hard sphere model it does not depend on the radius over the range shown. The results calculated using the Woods-Saxon models show only a slight variation. Part of this variation may be due to the difficulty in determining the second derivative from the numerical calculations. We recommend $[1/(2S)](d^2S/dE^2)|_{E=0} = 15.7 \text{ MeV}^{-2}$; the uncertainty in the second derivative introduces a negligible error in S_{eff} .

There has been considerable disagreement in the literature on what values should be used for the derivatives. (See for example, Ref. [5]). The last line of Table II gives values of the derivatives from Adelberger *et al.* [3] that are very different from those we have obtained in any of the other models. However, we can reproduce their results by following their procedure for determining the derivatives and fitting $S(E)$ by a quadratic form over the range $E = 20 - 300$ keV. Similarly, we can reproduce the values Barker obtained [5] for the derivatives by fitting a quadratic form over the range he chose, $0 - 100$ keV. An accurate determination of the derivatives from a fit of a quadratic form could only be obtained if the fit region was restricted to $E < 10$ keV. Thus the differences in the values

of derivatives quoted in the literature are due to the range of energies used for the fits and are unrelated to the true model dependence of the derivatives at threshold.

We show in Table III the results of using Eq. (5) with two different sets of derivatives. The first is that presented above, namely -1.87 MeV^{-1} and 15.7 MeV^{-2} , and the second is that from Adelberger *et al.* [3] given in Table II. Using this second set of derivatives introduces an error of 0.3 eVb in S_{eff} . As the reaction rate must be known to better than 5% [3], this 1.5% error is 30% of the total error allowed. Remarkably, using the inaccurate values of the derivatives at threshold is worse than neglecting the derivatives entirely and using $S(20)$.

In conclusion the use of approximate forms for S_{eff} does not generate significant errors if accurate parametrisations of the S factor and its derivatives are used. Eq. (7) is of special interest since it avoids the use of the derivatives by using S evaluated at 20 keV. We therefore recommend that determinations of the S factor for ${}^7\text{Be}(p, \gamma){}^8\text{B}$ at low energy quote $S(20)$ as well as $S(0)$.

ACKNOWLEDGMENTS

This work was motivated by discussions with C.W. Johnson. Financial support from Natural Sciences and Engineering Research Council of Canada is gratefully acknowledged.

REFERENCES

- [1] J. N. Bahcall and M. Pinsonneault, Rev. Mod. Phys. **67**, 885 (1992); J. N. Bahcall, S. Basu, and M.H. Pinsonneault, astro-ph/9805135.
- [2] J. N. Bahcall, Nucl. Phys. **75**, 10 (1966); J. N. Bahcall, Neutrino Astrophysics (Cambridge University, Cambridge).
- [3] E. Adelberger *et al.*, Rev. Mod. Phys (October, 1998, in press), astro-ph/9805121.
- [4] B. K. Jennings, S. Karataglidis, and T. D. Shoppa, nuth-ph/9806067, submitted to Phys. Rev. C.
- [5] F. C. Barker, Phys. Rev. C **28** 1407 (1983).

TABLES

TABLE I. The values of S_{eff} , in eVb, from the different approximations using Eq. (8) for $S(E)$. E_0 is in keV and T is in 10^6 K.

T	E_0	S_{eff}	$S_{\text{eff-S}}$	$S_{\text{eff-T}}$	$S_{\text{eff-Snd}}$	$S_{\text{eff-S20}}$
12	15.4	18.67	18.68	18.69	18.70	18.58
13	16.3	18.65	18.66	18.68	18.69	18.59
14	17.1	18.64	18.65	18.66	18.67	18.59
15	17.9	18.62	18.63	18.64	18.65	18.60
16	18.7	18.60	18.61	18.63	18.64	18.60
17	19.4	18.58	18.59	18.61	18.62	18.61

TABLE II. The first and second derivatives of S evaluated at threshold for the models of Ref. [4]. The nomenclature is also that of Ref. [4].

Model	$\frac{1}{S} \frac{dS}{dE}$ (MeV $^{-1}$)	$\frac{1}{2S} \frac{d^2S}{dE^2}$ (MeV $^{-2}$)
$r_c = 0.0$ fm	-1.77	15.7
$r_c = 1.0$ fm	-1.82	15.7
$r_c = 2.4$ fm	-1.92	15.7
$r_c = 4.1$ fm	-2.09	15.7
B1	-1.91	16.1
B2	-1.84	16.0
T	-1.96	15.7
Adelberger <i>et al.</i> [3]	-0.70	1.9

TABLE III. $S_{\text{eff-T}}$, in eVb, as obtained using the two different sets of derivatives discussed in the text. The temperature is in 10^6 K.

T	12	13	14	15	16	17
$S_{\text{eff-T}}$ (eVb)	18.68	18.66	18.64	18.62	18.61	18.59
$S_{\text{eff-T}}$ (eVb)	18.97	18.96	18.96	18.95	19.94	18.94